

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA  
FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA  
ESCUELA DE MEDICINA VETERINARIA**



**ANTIBIOTIC RESISTANCE AND SENSITIVITY  
PATTERNS OF BACTERIA ISOLATED FROM MASTITIS  
CASES IN DAIRY COWS IN HONDURAS**

**RESISTENCIA ANTIBIÓTICA Y SENSIBILIDAD EN  
AISLAMIENTO EN BACTERIAS EN MASTITIS EN  
VACAS LECHERAS EN HONDURAS**

**Médica Veterinaria**

**GUATEMALA, MAYO DE 2021**

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA  
FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA  
ESCUELA DE MEDICINA VETERINARIA**



**ANTIBIOTIC RESISTANCE AND SENSITIVITY  
PATTERNS OF BACTERIA ISOLATED FROM MASTITIS  
CASES IN DAIRY COWS IN HONDURAS**

**RESISTENCIA ANTIBIÓTICA Y SENSIBILIDAD EN  
AISLAMIENTO EN BACTERIAS EN MASTITIS EN  
VACAS LECHERAS EN HONDURAS**

**TRABAJO DE GRADUACIÓN**

**PRESENTADO A LA HONORABLE JUNTA DIRECTIVA DE LA FACULTAD**

**POR**

**DANIELA GEOVANNA MARTÍNEZ MATAMOROS**

Al conferírsele el título profesional

**Médica Veterinaria**

En el grado de Licenciado

**GUATEMALA, MAYO DE 2021**

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA  
FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA  
ESCUELA DE MEDICINA VETERINARIA**

DECANO:	M.A. Rodolfo Chang Shum
SECRETARIO:	M. Sc. Lucrecia Emperatriz Motta Rodriguez
VOCAL I:	M. Sc. Juan José Prem González
VOCAL II:	Lic. Zoot. Miguel Ángel Rodenas Argueta
VOCAL III:	M.V. Edwin Rigoberto Herrera Villatoro
VOCAL IV:	P. Agr. Luis Gerardo López Morales
VOCAL V:	Br. María José Solares Herrera

**ASESORES**

PHD. DENNIS SIGFRIED GUERRA CENTENO  
Dr. Sc. CARLOS VALDEZ SANDOVAL  
M. Sc. MANUEL LEPE LÓPEZ

En cumplimiento con lo establecido por los reglamentos y normas de la Universidad de San Carlos de Guatemala, presento a su consideración el artículo científico publicado en una revista indexada, titulado:

**ANTIBIOTIC RESISTANCE AND SENSITIVITY  
PATTERNS OF BACTERIA ISOLATED FROM MASTITIS  
CASES IN DAIRY COWS IN HONDURAS**

**RESISTENCIA ANTIBIÓTICA Y SENSIBILIDAD EN  
AISLAMIENTO EN BACTERIAS EN MASTITIS EN  
VACAS LECHERAS EN HONDURAS**

Que fuera aprobado por la Honorable Junta Directiva de la Facultad de  
Medicina Veterinaria y Zootecnia

Como requisito previo a optar el título de:

**Médica Veterinaria**

## **ACTO QUE DEDICO A:**

### **A Dios:**

Por permitirme alcanzar uno de mis sueños culminando este capítulo. Porque siempre me guio en cada momento de mi vida.

### **A mis padres:**

Rosa Maria y Geovanni, por enseñarme los valores y darme su amor incondicional siempre. Por qué nunca dejaron de creer en mí y siempre dieron todo para que pudiera cumplir con esta meta. Este logro es nuestro.

## **AGRADECIMIENTOS**

- A Dios:** Por todas las bendiciones que he recibido, una familia que siempre me ha apoyado y sabiduría para llegar hasta aquí.
- A mis padres:** Gracias por todo este sacrificio, esfuerzo y paciencia de tener que estar aparte durante mi ciclo universitario. Por eso y mucho más nunca dejare de estar agradecida.
- A mis hermanos:** A David y Kevin, por el apoyo moral y emocional en mi vida. Por el esfuerzo de madrugada y noches siempre que volvía a Honduras.
- A mi abuela:** Por brindar sus consejos y motivaciones en cada momento de mi vida.
- A mis tios:** Por enseñarme que lo más importante es la unión familiar y sus consejos de vida.
- A mis asesores:** Por darme la oportunidad de trabajar con grandes profesionales en la Medicina Veterinaria.
- A todos mis amigos:** Gracias por todos esos momentos inolvidables que tuvimos y por hacerme sentir en casa en un país extranjero.
- Al Dr. Sergio Véliz:** Gracias por ser un gran amigo y colega; gracias por compartir sus consejos y conocimientos en todo momento.

# ÍNDICE

I.	SUMMARY.....	1
II.	RESUMEN.....	1
III.	INTRODUCTION.....	1
IV.	MATERIAL AND METHODS.....	2
V.	RESULTS.....	2
VI.	TABLE I.....	3
VII.	TABLE II.....	4
VIII.	TABLE III.....	5
IX.	TABLE IV.....	6
X.	TABLE V.....	7
XI.	DISCUSSION.....	2,8
XII.	CONCLUSION.....	8
XIII.	ACKNOWLEDGMENT.....	8
XIV.	BIBLIOGRAGPY.....	8,9,10



## Antibiotic resistance and sensitivity patterns of bacteria isolated from mastitis cases in dairy cows in Honduras

Martínez-Matamoros, D. <sup>1,2</sup>; Guerra-Centeno, D. <sup>2@</sup>; Lepe-López, M. <sup>2</sup>; and Valdez-Sandoval, C. <sup>2</sup>

<sup>1</sup>Servicio Nacional de Sanidad e Inocuidad Agroalimentaria (SENASA), Honduras.

<sup>2</sup>Instituto de Investigación en Ciencia Animal y Ecosalud, Facultad de Medicina Veterinaria y Zootecnia, Universidad de San Carlos de Guatemala. Guatemala.

### SUMMARY

Mastitis is a major problem in dairy farms around the world and is usually treated with antibiotics. However, bacteria - the usual causative agents - often become resistant to commonly used antibiotics, hindering milk production, and generating a possible public health problem. With the purpose of generating information on this situation in Honduras, we carried out a retrospective documentary study examining official records of bacterial isolates and antibiotic resistance tests carried out to milk samples coming from mastitis cases in dairy cows from 2013 to 2018. We examined 235 cases of bacterial isolation and sensitivity test to 15 antibiotics (amoxicillin + clavulanic acid, ampicillin, ceftiofur, ceftriaxone, enrofloxacin, ciprofloxacin, oxacillin, doxycycline, neomycin, trimethoprim + sulphamethoxazole, gentamicin, tetracycline, erythromycin, and penicillin). The more frequently isolated bacteria taxon was *Staphylococcus* spp. (168 cases) followed by *Streptococcus* spp. (31 cases), *Corynebacterium* spp. (14 cases), *Escherichia* spp. (13 cases), *Klebsiella* spp. (five cases), *Enterobacter* sp. (one case), *Pasteurella* sp. (one case), and *Pseudomonas* sp. (one case). We found antibiotic resistance cases to practically all tested antibiotics, however, patterns in resistance and sensitivity were uncommon.

### Resistencia antibiótica y sensibilidad en aislamientos de bacterias en mastitis en vacas lecheras en Honduras

### RESUMEN

La mastitis es un importante problema en granjas de producción lechera alrededor del mundo y usualmente se trata con antibióticos. Sin embargo, las bacterias -que son los agentes causales usuales- van volviéndose resistentes a los antibióticos de uso común. Con el objetivo de generar información sobre esta situación en Honduras, realizamos un estudio retrospectivo documental examinando los registros oficiales de aislamientos de bacterias y pruebas de resistencia antibiótica llevados a cabo en muestras de leche de casos de mastitis en vacas, entre 2013 y 2018. Examinamos 235 casos de aislamiento bacteriano y pruebas de sensibilidad a 15 antibióticos (amoxicilina + ácido clavulánico, ampicilina, ceftiofur, ceftriaxona, enrofloxacina, ciprofloxacina, oxacilina, doxiciclina, neomicina, trimetoprim + sulfametoxazol, gentamicina, tetraciclina, eritromicina y penicilina). Los taxones más frecuentemente aislados fueron *Staphylococcus* spp. (168 casos), seguido por *Streptococcus* spp. (31 casos), *Corynebacterium* spp. (14 casos), *Escherichia coli* (13 casos), *Klebsiella* spp. (cinco casos), *Enterobacter* sp. (un caso), *Pasteurella* sp. (un caso) y *Pseudomonas* sp. (un caso). Encontramos casos de resistencia a prácticamente todos los antibióticos, aunque los patrones de resistencia fueron poco comunes.

### AdditionAl keywords

Antimicrobial resistance.  
Mastitis therapy.  
Milk bacteria.  
Udder health

### PAIAbrAs clAve

Bacterias de la leche.  
Resistencia antimicrobiana.  
Salud de la ubre.  
Terapia de mastitis.

### InformAtion

Cronología del artículo.  
Recibido/Received: 06.05.2020  
Aceptado/Accepted: 02.11.2020  
On-line: 15.01.2021  
Correspondencia a los autores/Contact e-mail:  
[m.sc.dennisguerra@gmail.com](mailto:m.sc.dennisguerra@gmail.com)

### INTRODUCTION

The worldwide growth of bacterial resistance to antibiotics has been identified as a major public health

threat (Roca et al., 2015; World Health Organization, 2014). Animal production has been considered one of the main sources of resistant or multi-resistant bacterial strains (Angulo et al., 2014; Eng, 2015; Ventola, 2015).

The massive and routine use of antibiotics in animal production, either for prevention or therapy, exerts a selection pressure over

bacterial populations thus promoting the appearance of super-resistant strains (Aarestrup, 2012; Dahms et al., 2014). It has been estimated that the global consumption of antibiotics in animal production will increase by 67% between 2010 and 2030 (Van Boeckel et al., 2015) and this, of course, could aggravate the problem.

Cow milk production is one of the most common farm activities around the world (Faye & Konuspaeva, 2012) and mastitis is one of the most common



*Arch. Zootec. 70 (269): 50-59. 2021.*

diseases and certainly a frequent reason for anti-biotic use (Barlow, 2011). This means that not only could the milk of a cow with mastitis harbor bacteria, but these bacteria could be antibiotic-resistant. In a developing country like Honduras, where the government is not able to control all the milk or dairy products that are sold on the markets, there is an increased risk of acquiring antibiotic-resistant bacteria via the consumption of these products.

Despite Honduras is a traditional dairy farmer country (Molina, 2010), the publications about mastitis-causing bacteria and their antibiotic-resistant patterns are practically inexistent. Therefore, our objectives in this research were to analyze official records of laboratory culture and sensitivity tests of mastitis cases in dairy cows in Honduras from 2013 to 2018.

## MATERIAL AND METHODS

### Study Site and Sampling

We conducted a retrospective study based on records of the Honduran Institute of Veterinary Medical Research (HIVMR), National Service of Agrifood Health and Safety, Republic of Honduras. We examined the outcomes of microbiologic cultures of milk samples from mastitis cases in dairy cows, from years 2013 to 2018. We looked for bacterial isolation and antibiotic sensitivity tests. The samples came from 12 Honduran provinces (Atlántida, Choluteca, Colón, Comayagua, Cortés, El Paraíso, Francisco Morazán, Intibucá, La Paz, Olancho, Santa Bárbara, Yoro).

The isolation and identification of bacteria in the HIVMR was carried out according to the ISO 6579 International Standard Procedures (International Organization for Standardization, 2002) and the antimicrobial susceptibility tests were performed following the methods and criteria of the M100S standards of the Clinical and Laboratory Standards Institute (Patel et al., 2017).

### Data processing

Positive cases were examined and the isolated species (or taxon) of bacteria was registered as well as the sensitivity to 15 common use antibiotics (Amoxicillin + Clavulanic Acid, Ampicillin, Ceftiofur, Ceftriaxone, Enrofloxacin, Ciprofloxacin, Oxacillin, Doxycycline, Neomycin, Trimethoprim + Sulphamethoxazole, Gentamicin, Tetracycline, Erythromycin, and Penicillin). We arranged the data by the species of bacteria isolated, province of origin, year of sample processing and sensitivity test outcomes.

### Statistical analysis

We looked for patterns in isolations as well as antibiotic resistance and sensitivity throughout the

six years of data using heterogeneity G-tests (Sokal & Rohlf, 1995). All tests were performed in the software R. Tests were not performed for periods that had less than two years of data.

## RESULTS

We examined 263 mastitis clinical case records and found 235 positives to bacterial isolation and sensitivity test. Thirteen cases corresponded to Atlántida, seven to Choluteca, 13 to Colón, 11 to Comayagua, 15 to Cortés, 19 to El Paraíso, 78 to Francisco Morazán, three to Intibucá, 40 to La Paz, 17 to Olancho, 16 to Santa Bárbara and three to Yoro.

The more frequently isolated bacteria taxon was *Staphylococcus* spp. (168 cases) followed by *Streptococcus* spp. (31 cases), *Corynebacterium* spp. (14 cases), *Escherichia* spp. (13 cases), *Klebsiella* sp. (five cases), *Enterobacter* sp. (one case), *Pasteurella* sp. (one case) and *Pseudomonas* sp. (one case). We found antibiotic resistance cases to practically all tested antibiotics. The sensitivity and resistance patterns shown by the four most isolated taxa to 15 common use antibiotics from 2013 to 2018 are shown in tables I to IV. The summation of resistance and sensitivity frequencies for these taxa is shown in Table V.

In general, patterns in resistance or sensitivity were infrequent. However, for the most isolated taxon (*Staphylococcus* spp.), we observed just a few patterns and these patterns did not show any strong tendency towards sensitivity or resistance (Table I). For the least isolated taxa, the observed patterns are probably not conclusive.

## DISCUSSION

The genus *Staphylococcus* is a common etiologic agent of intramammary infection in cows (Sutra & Poutrel, 1994). Several species have been identified in clinical and subclinical cases (Hosseinzadeh & Dastmalchi-Saei, 2014; Rall et al., 2014). Their presence is ubiquitous in the environment, as well as the skin and mucous membranes of cows, so it can easily invade the nipple and the mammary gland (DeGo et al., 2002). According to our data, *Staphylococcus* spp. were involved in 168 of 234 mastitis records. Furthermore, 685 of the 1836 antibiotic sensitivity tests performed for this taxon yielded resistance outcomes.

Although our data are probably not representative of the Honduran dairy cow population, they do suggest that antibiotic treatment of mastitis caused by *Staphylococcus* spp. could fail 37% of the time. Antibiotic resistance of *Staphylococcus* spp. has been previously reported in Algeria (Saidi et al., 2019), Argentina (Gentilini et al., 2000), Brazil (Coelho et al., 2009), Germany (Lüthje & Schwarz, 2006), Iran (Jamali et al., 2014), Korea (Moon et al., 2007), Switzerland (Frey et al., 2013) and Turkey (Güler et al.,



**Table I. Antibiotic sensitivity and resistance frequencies of *Staphylococcus* spp. isolated from bovine mastitis cases submitted to the Honduran Institute of Medical Veterinary Research (IHIMV), 2013 to 2018 (Sensibilidad frente antibióticos y frecuencias de resistencia de *Staphylococcus* spp. aisladas de casos de mastitis bovina presentadas al Instituto Hondureño de Investigaciones Médico Veterinarias (IHIMV), 2013 a 2018).**

No.	Antibiotic		2013	2014	2015	2016	2017	2018	Total
1	Amoxicillin + Clavulanic Acid**	S	7	12	0	6	26	20	72
		IS	0	0	0	0	0	0	0
		R	2	13	11	3	9	4	42
2	Ampicillin**	S	6	8	0	0	14	21	49
		IS	0	0	0	0	0	0	0
		R	20	36	11	0	22	9	100
3	Ceftiofur**	S	14	8	7	5	19	27	82
		IS	0	10	4	3	0	3	20
		R	7	18	0	2	7	0	34
4	Ceftriaxone**	S	16	20	0	0	12	20	69
		IS	0	16	11	0	0	4	31
		R	8	6	0	0	6	2	22
5	Cefuroxime <sup>□</sup>	S	0	5	0	0	0	0	5
		IS	0	9	0	0	0	1	10
		R	0	0	0	0	0	0	0
6	Enrofloxacin*	S	13	24	5	10	23	11	87
		IS	3	3	0	0	0	3	9
		R	4	2	0	0	0	2	8
7	Ciprofloxacin <sup>□</sup>	S	16	15	0	0	14	20	66
		IS	2	2	0	0	1	1	6
		R	8	15	0	0	4	4	31
8	Oxacillin**	S	6	20	7	7	19	19	80
		IS	0	2	0	0	0	0	2
		R	17	21	0	3	12	7	60
9	Doxycycline**	S	2	13	3	7	14	13	53
		IS	0	3	8	0	1	0	13
		R	3	8	0	3	10	3	27
10	Neomycin**	S	7	16	0	4	14	10	51
		IS	0	9	7	4	2	0	23
		R	15	17	0	0	5	5	42
11	Trimethoprim + Sulfamethoxazole**	S	17	27	11	9	23	17	106
		IS	0	5	0	0	0	0	5
		R	8	6	0	1	8	10	33
12	Gentamicin**	S	16	29	3	10	29	27	114
		IS	1	1	0	0	0	0	2
		R	10	20	4	0	1	3	38
13	Tetracycline <sup>□</sup>	S	14	20	3	5	18	13	74
		IS	0	7	0	2	4	4	17
		R	10	22	8	3	10	6	60
14	Erythromycin**	S	6	17	0	1	7	16	48
		IS	3	7	11	4	2	8	36
		R	17	23	0	1	8	3	52
15	Penicillin**	S	4	11	0	3	13	17	48
		IS	0	4	0	0	2	0	6
		R	23	34	11	7	19	7	103

Note: S=sensitive, IS=intermediate sensitivity, R=resistant, <sup>□</sup>=pattern (homogeneity), \*=significant heterogeneity, \*\*=highly significant heterogeneity.



**Table II.** Antibiotic sensitivity and resistance frequencies of *Streptococcus* spp. isolated from bovine mastitis cases submitted to the Honduran Institute of Medical Veterinary Research (IHIMV), 2013 to 2018 (Sensibilidad frente antibióticos y frecuencias de resistencia de *Streptococcus* spp. aisladas de casos de mastitis bovina presentadas al Instituto Hondureño de Investigaciones Médico Veterinarias (IHIMV), 2013 a 2018).

No.	Antibiotic		2013	2014	2015	2016	2017	2018	Total
1	Amoxicillin + Clavulanic Acid <sup>□</sup>	S	8	3	---	1	5	5	22
		IS	0	0	---	0	0	0	0
		R	3	0	---	2	0	0	5
2	Ampicillin <sup>□</sup>	S	6	1	---	1	5	1	14
		IS	1	0	---	0	1	3	5
		R	4	2	---	0	0	2	8
3	Ceftiofur <sup>□</sup>	S	7	1	---	2	3	6	19
		IS	2	0	---	0	0	0	2
		R	0	2	---	0	0	0	2
4	Ceftriaxone**	S	11	1	---	0	4	0	16
		IS	0	4	---	0	0	3	7
		R	0	0	---	0	0	1	1
5	Cefuroxime <sup>□</sup>	S	6	1	---	0	0	0	7
		IS	2	0	---	0	0	2	4
		R	3	0	---	0	0	0	3
6	Enrofloxacin <sup>□</sup>	S	7	0	---	2	2	0	11
		IS	2	0	---	1	1	0	4
		R	2	2	---	0	0	0	4
7	Ciprofloxacin*	S	5	1	---	0	3	0	9
		IS	2	0	---	0	0	3	5
		R	4	2	---	0	0	0	6
8	Oxacillin*	S	9	1	---	0	5	0	15
		IS	0	0	---	0	0	0	0
		R	2	2	---	3	1	5	13
9	Doxycycline <sup>□</sup>	S	5	1	---	1	3	2	12
		IS	0	0	---	0	0	0	0
		R	4	2	---	2	0	3	11
10	Neomycin <sup>□</sup>	S	0	1	---	0	3	0	4
		IS	0	2	---	0	0	0	2
		R	0	2	---	1	1	5	9
11	Trimethoprim + Sulfamethoxazole**	S	4	0	---	0	5	0	9
		IS	2	0	---	0	1	0	3
		R	5	4	---	3	0	6	18
12	Gentamicin**	S	11	1	---	2	5	0	19
		IS	0	0	---	0	0	1	1
		R	0	2	---	1	1	5	9
13	Tetracycline**	S	5	3	---	0	0	2	10
		IS	0	1	---	0	5	0	6
		R	6	1	---	3	1	4	15
14	Erythromycin <sup>□</sup>	S	4	2	---	1	3	3	13
		IS	2	1	---	0	0	2	5
		R	5	2	---	2	0	1	10
15	Penicillin**	S	3	3	---	0	5	1	12
		IS	4	0	---	0	0	0	4
		R	4	2	---	3	0	5	14

Note: S = sensitive, IS = intermediate sensitive, R = resistant, <sup>□</sup> = pattern (homogeneity), \* = significant heterogeneity, \*\* = highly significant heterogeneity .



**Table III. Antibiotic sensitivity and resistance frequencies of *Corynebacterium* sp. isolated from bovine mastitis cases submitted to the Honduran Institute of Medical Veterinary Research (IHIMV), 2013 to 2018** (Sensibilidad a los antibióticos y frecuencias de resistencia de *Corynebacterium* sp. aisladas de casos de mastitis bovina presentadas al Instituto Hondureño de Investigaciones Médico Veterinarias (IHIMV), 2013 a 2018).

No.	Antibiotic		2013	2014	2015	2016	2017	2018	Total
1	Amoxicillin + Clavulanic Acid**	S	1	---	---	0	0	20	4
		IS	0	---	---	0	0	0	0
		R	0	---	---	6	0	4	0
2	Ampicillin <sup>□</sup>	S	1	---	---	0	0	21	3
		IS	0	---	---	0	0	0	0
		R	0	---	---	0	0	9	2
3	Ceftiofur**	S	0	---	---	0	0	27	5
		IS	0	---	---	4	0	3	0
		R	0	---	---	2	0	0	0
4	Ceftriaxone <sup>□</sup>	S	1	---	---	0	0	20	4
		IS	0	---	---	0	0	4	0
		R	0	---	---	0	0	2	0
5	Cefuroxime <sup>□</sup>	S	0	---	---	0	0	0	0
		IS	0	---	---	0	0	1	0
		R	0	---	---	0	0	0	0
6	Enrofloxacin <sup>□</sup>	S	0	---	---	4	2	11	4
		IS	1	---	---	2	0	3	0
		R	0	---	---	0	0	2	0
7	Ciprofloxacin <sup>□</sup>	S	1	---	---	0	0	20	4
		IS	0	---	---	0	0	1	0
		R	0	---	---	0	0	4	0
8	Oxacillin*	S	0	---	---	0	0	19	0
		IS	0	---	---	0	0	0	0
		R	0	---	---	4	2	7	2
9	Doxycycline*	S	0	---	---	0	0	13	3
		IS	0	---	---	0	0	0	0
		R	0	---	---	6	0	3	0
10	Neomycin <sup>□</sup>	S	0	---	---	2	2	10	2
		IS	0	---	---	0	0	0	1
		R	0	---	---	0	0	5	0
11	Trimethoprim + Sulfamethoxazole <sup>□</sup>	S	0	---	---	4	0	17	4
		IS	0	---	---	0	0	0	0
		R	1	---	---	2	2	10	1
12	Gentamicin**	S	11	1	---	2	5	0	19
		IS	0	0	---	0	0	1	1
		R	0	2	---	1	1	5	9
13	Tetracycline**	S	5	3	---	0	0	2	10
		IS	0	1	---	0	5	0	6
		R	6	1	---	3	1	4	15
14	Erythromycin <sup>□</sup>	S	4	2	---	1	3	3	13
		IS	2	1	---	0	0	2	5
		R	5	2	---	2	0	1	10
15	Penicillin**	S	3	3	---	0	5	1	12
		IS	4	0	---	0	0	0	4
		R	4	2	---	3	0	5	14

Note: S=sensitive, IS=intermediate sensitivity, R=resistant, <sup>□</sup>=pattern (homogeneity), \*=significant heterogeneity, \*\*=highly significant heterogeneity.



**Table IV. Antibiotic sensitivity and resistance frequencies of *Escherichia* spp. isolated from bovine mastitis cases submitted to the Honduran Institute of Medical Veterinary Research (IHIMV), 2013 to 2018** (Sensibilidad a los antibióticos y frecuencias de resistencia de *Escherichia* spp. aisladas de casos de mastitis bovina presentadas al Instituto Hondureño de Investigaciones Médico Veterinarias (IHIMV), 2013 a 2018).

No.	Antibiotic		2013	2014	2015	2016	2017	2018	Total
1	Amoxicillin + Clavulanic Acid <sup>□</sup>	S	5	0	---	---	1	---	6
		IS	0	0	---	---	0	---	0
		R	0	1	---	---	0	---	1
2	Ampicillin*	S	0	3	---	---	1	---	4
		IS	0	0	---	---	0	---	0
		R	5	2	---	---	0	---	7
3	Ceftiofur <sup>□</sup>	S	0	0	---	---	0	---	0
		IS	0	3	---	---	0	---	3
		R	0	0	---	---	0	---	0
4	Ceftriaxone <sup>□</sup>	S	5	5	---	---	1	---	11
		IS	0	0	---	---	0	---	0
		R	0	2	---	---	0	---	2
5	Cefuroxime <sup>□</sup>	S	2	3	---	---	0	---	5
		IS	0	0	---	---	0	---	0
		R	0	0	---	---	0	---	0
6	Enrofloxacin <sup>□</sup>	S	5	2	---	---	0	---	7
		IS	0	0	---	---	0	---	0
		R	0	0	---	---	0	---	0
7	Ciprofloxacin <sup>□</sup>	S	2	3	---	---	1	---	6
		IS	0	0	---	---	0	---	0
		R	0	2	---	---	0	---	2
8	Oxacillin <sup>□</sup>	S	5	1	---	---	0	---	6
		IS	0	0	---	---	0	---	0
		R	0	2	---	---	0	---	2
9	Doxycycline*	S	2	3	---	---	0	---	5
		IS	0	4	---	---	0	---	4
		R	3	0	---	---	0	---	3
10	Neomycin <sup>□</sup>	S	0	2	---	---	0	---	2
		IS	2	5	---	---	1	---	8
		R	0	0	---	---	0	---	0
11	Trimethoprim + Sulfamethoxazole**	S	0	4	---	---	1	---	5
		IS	2	0	---	---	0	---	2
		R	3	0	---	---	0	---	3
12	Gentamicin <sup>□</sup>	S	5	3	---	---	1	---	9
		IS	0	0	---	---	0	---	0
		R	0	0	---	---	0	---	0
13	Tetracycline <sup>□</sup>	S	0	5	---	---	0	---	5
		IS	0	0	---	---	0	---	0
		R	5	2	---	---	1	---	8
14	Erythromycin <sup>□</sup>	S	0	3	---	---	0	---	3
		IS	0	0	---	---	0	---	0
		R	0	4	---	---	0	---	4
15	Penicillin <sup>□</sup>	S	0	1	---	---	0	---	1
		IS	0	0	---	---	0	---	0
		R	0	2	---	---	0	---	2

Note: S = sensitive, IS = intermediate sensitivity, R = resistant, <sup>□</sup> = pattern (homogeneity), \* = significant heterogeneity, \*\* = highly significant heterogeneity.



**Table V. Antibiotic sensitivity and resistance number of outcomes for the more frequently isolated taxa from bovine mastitis cases submitted to the Honduran Institute of Medical Veterinary Research (IHIMV), 2013 to 2018**

(Sensibilidad a los antibióticos y número de resistencia de los resultados para los taxones más frecuentemente aislados de los casos de mastitis bovina presentados al Instituto Hondureño de Investigaciones Médico Veterinarias (IHIMV), 2013 a 2018).

Antibiotic	Challenge outcome	Taxon			
		<i>Staphylococcus</i>	<i>Streptococcus</i>	<i>Corynebacterium</i>	<i>Escherichia</i>
Amoxicillin + Clavulanic Acid	S	72	22	5	6
	IS	0	0	0	0
	R	42	5	6	1
Ampicillin	S	49	14	4	4
	IS	0	5	0	0
	R	100	8	2	7
Ceftiofur	S	82	19	5	0
	IS	20	2	4	3
	R	34	2	2	0
Ceftriaxone	S	69	16	5	11
	IS	31	7	0	0
	R	22	1	0	2
Cefuroxime	S	5	7	0	5
	IS	10	4	0	7
	R	0	3	0	0
Enrofloxacin	S	87	11	10	7
	IS	9	4	3	0
	R	8	4	0	0
Ciprofloxacin	S	66	9	5	6
	IS	6	5	0	0
	R	31	6	0	2
Oxacillin	S	80	15	0	6
	IS	2	0	0	0
	R	60	13	8	2
Doxycycline	S	53	12	3	5
	IS	13	0	0	4
	R	27	11	6	3
Neomycin	S	51	4	6	2
	IS	23	2	1	8
	R	42	9	0	0
Trimethoprim + Sulfamethoxazole	S	106	9	8	5
	IS	5	3	0	2
	R	33	18	6	3
Gentamicin	S	114	19	8	9
	IS	2	1	1	0
	R	38	9	3	0
Tetracycline	S	74	10	2	5
	IS	17	6	1	0
	R	60	15	8	8
Erythromycin	S	48	13	1	3
	IS	36	5	0	0
	R	52	10	9	4
Penicillim	S	48	12	1	1
	IS	6	4	1	0

Note: S = sensitive, IS = intermediate sensitivity, R = resistant.



2005) among others. The high frequencies of resistance showed by this taxon in this study to ampicillin, ciprofloxacin, oxacillin, doxycycline, neomycin, and tetracycline is noteworthy, and had not been previously reported.

*Streptococcus* spp., the second most frequently isolated taxon in this study, is also one of the common causative agents of mastitis in cows (Oikonomou et al., 2012). Three species (*S. agalactiae*, *S. dysgalactiae*, and *S. uberis*) have been described as the most frequently isolated (Baseggio et al., 1997). As for antibiotic resistance, there are reports in China (Gao et al., 2012), Egypt (El-Jakee et al., 2013), France (Guérin-Faubleé et al., 2002), Germany (Minst et al., 2012), New Zealand (McDougall et al., 2013), Switzerland (Corti et al., 2003) and others, although none of these have reported the high frequencies of resistance to enrofloxacin, ciprofloxacin, oxacillin, doxycycline, neomycin, trimethoprim + sulfamethoxazole, and erythromycin seen in our study.

*Corynebacterium* spp. and *Escherichia coli* were less frequent in our study. Nevertheless, the resistance to oxacillin, doxycycline, trimethoprim + sulfamethoxazole, tetracycline, erythromycin, and penicillin, and the sensitivity to enrofloxacin in both taxa was evident.

The high frequencies of antibiotic resistance observed in this study, in contrast to the low frequencies observed in other countries like Finland (Pitkälä et al., 2004), France (Botrel et al., 2010), Portugal (Rato et al., 2013) may be due to cultural differences in mastitis management practices. The improper and uncontrolled use of antibiotics has probably contributed to the observed resistance of mastitis-causing bacteria. Dairy farmers in Honduras, face a rising challenge to successfully treat mastitis. Furthermore, the massive and routine use of antibiotics and the proliferation of resistant bacteria strains could lead to unknown impacts on dairy product consumers. Bando et al. (2009), found over 41% prevalence of antibiotic residues in pasteurized milk samples in Paraná, Brazil and this could be a reality in other countries where mastitis caused by antibiotic-resistant bacteria is common.

## CONCLUSIONS

The more frequently isolated taxa were *Staphylococcus* spp., followed by *Streptococcus* spp., *Corynebacterium* spp., and *Escherichia* spp. We found resistance cases to practically all the tested antibiotics (even the broad-spectrum ones), however, resistance patterns were not common throughout the evaluated period. More research needs to be done to determine the mastitis-causing bacteria in Honduras to the species level and to better understand the ecological relationship between bacterial species and dairy cows.

## ACKNOWLEDGMENTS

We want to thank Dr. Rafael Rodríguez, head of the epidemiology department of the National Service of Agri-food Health and Safety of Honduras (SENASA) for the access to laboratory records.

## BIBLIOGRAPHY

- Aarestrup, F 2012, 'Get pigs off antibiotics', *Nature*, vol. 486, no. 7404, pp. 465-466. <https://doi.org/10.1038/486465a>
- Angulo, F J, Baker, N L, Olsen, S J, Anderson, A, & Barrett, T J 2004, 'Antimicrobial use in agriculture: controlling the transfer of antimicrobial resistance to humans', *Seminars in pediatric infectious diseases*, vol. 15, no. 2, pp. 78-85. <https://doi.org/10.1053/j.spid.2004.01.010>
- Bando, E, Oliveira, R C, Ferreira, G M Z, & Machinski, J M 2009, 'Occurrence of antimicrobial residues in pasteurized milk commercialized in the state of Paraná, Brazil', *Journal of food protection*, vol. 72 no. 4, pp. 911-914. <https://doi.org/10.4315/0362-028X-72.4.911>
- Barlow, J 2011, 'Mastitis therapy and antimicrobial susceptibility: a multispecies review with a focus on antibiotic treatment of mastitis in dairy cattle' *Journal of mammary gland biology and neoplasia*, vol. 16, no. 4, pp. 383-407. <https://doi.org/10.1007/s10911-011-9235-z>
- Baseggio, N, Mansell, P D, Browning, J W, & Browning, G F 1997, 'Strain differentiation of isolates of streptococci from bovine mastitis by pulsed-field gel electrophoresis' *Molecular and cellular probes*, vol. 11, no. 5, pp. 349-354. <https://doi.org/10.1006/mcpr.1997.0126>
- Botrel, M A, Haenni, M, Morignat, E, Sulpice, P, Madec, J Y, & Calavas, D 2010, 'Distribution and antimicrobial resistance of clinical and subclinical mastitis pathogens in dairy cows in Rhône-Alpes, France' *Foodborne pathogens and disease*, vol. 7, no. 5, pp. 479-487. <https://doi.org/10.1089/fpd.2009.0425>
- Coelho, S M, Reinoso, E, Pereira, I A, Soares, L C, Demo, M, Bogni, C, & Souza, M 2009, 'Virulence factors and antimicrobial resistance of *Staphylococcus aureus* isolated from bovine mastitis in Rio de Janeiro' *Pesquisa Veterinária Brasileira*, vol. 29, no. 5, pp. 369-374. <https://doi.org/10.1590/S0100-736X2009000500002>
- Corti, S, Sicher, D, Regli, W, & Stephan, R 2003, 'Aktuelle Daten zur Antibiotikaresistenz der wichtigsten bovinen Mastitiserreger in der Schweiz' *Schweizer Archiv für Tierheilkunde*, vol. 14, no. 12, pp. 571-575. <https://doi.org/10.1024//0036-7281.145.12.571>
- Dahms, C, Hübner, N O, Wilke, F, & Kramer, A 2014, 'Mini-review: Epidemiology and zoonotic potential of multiresistant bacteria and *Clostridium difficile* in livestock and food' *GMS hygiene and infection control*, vol. 9, no. 3, pp. 1-16. <https://doi.org/10.3205/dgkh000241>
- DeGo, O K, Van Dijk, J E, & Nederbragt, H 2002, 'Factors involved in the early pathogenesis of bovine *Staphylococcus aureus* mastitis with emphasis on bacterial adhesion and invasion. A review' *Veterinary Quarterly*, vol. 24, no. 4, pp. 181-198. <https://doi.org/10.1080/01652176.2002.9695135>
- El-Jakee, J, Hableel, H S, Kandil, M, Hassan, O F, Khairy, E A, & Marouf, S A 2013, 'Antibiotic resistance patterns of isolated from mastitic cows and ewes in Egypt' *Global Veterinaria*, vol. 10, no. 3, pp. 264-270. <https://doi.org/10.5829/idosi.gv.2013.10.3.7248>



- Eng, S K, Pusparajah, P, Ab Mutalib, N S, Ser, H L, Chan, K G, & Lee, L H 2015, 'Salmonella: a review on pathogenesis, epidemiology and antibiotic resistance' *Frontiers in Life Science*, vol. 8, no. 3, pp. 284-293. <https://doi.org/10.1080/21553769.2015.1051243>
- Faye, B, & Konuspayeva, G 2012, 'The sustainability challenge to the dairy sector—The growing importance of non-cattle milk production worldwide' *International dairy journal*, vol. 24, no. 2, pp. 50-56. <https://doi.org/10.1016/j.idairyj.2011.12.011>
- Frey, Y, Rodriguez, J P, Thomann, A, Schwendener, S, & Perreten, V 2013, 'Genetic characterization of antimicrobial resistance in coagulase-negative staphylococci from bovine mastitis milk' *Journal of dairy science*, vol. 96, no. 4, pp. 2247-2257. <https://doi.org/10.3168/jds.2012-6091>
- Gao, J, Yu, F Q, Luo, L P, He, J Z, Hou, R G, Zhang, H Q, ... & Han, B 2012, 'Antibiotic resistance of Streptococcus agalactiae from cows with mastitis' *The Veterinary Journal*, vol. 194, no. 3, pp. 423-424. <https://doi.org/10.1016/j.tvjl.2012.04.020>
- Gentilini, E, Denamiel, G, Betancor, A, Rebuelto, M, Fermepin, M R, & De Torres, R A 2002, 'Antimicrobial susceptibility of coagulase-negative staphylococci isolated from bovine mastitis in Argentina' *Journal of Dairy Science*, vol. 85, no. 8, pp. 1913-1917. [https://doi.org/10.3168/jds.S0022-0322\(00\)74988-5](https://doi.org/10.3168/jds.S0022-0322(00)74988-5)
- Guérin-Faubleé, V, Tardy, F, Bouveron, C, & Carret, G 2002, 'Antimicrobial susceptibility of Streptococcus species isolated from clinical mastitis in dairy cows' *International journal of antimicrobial agents*, vol. 19, no. 3, pp. 219-226. [https://doi.org/10.1016/S0924-8579\(01\)00485-X](https://doi.org/10.1016/S0924-8579(01)00485-X)
- Güler, L, Ok, Ü, Gündüz, K, Gülcü, Y, & Hadimli, H H 2005, 'Antimicrobial susceptibility and coagulase gene typing of Staphylococcus aureus isolated from bovine clinical mastitis cases in Turkey' *Journal of dairy science*, vol. 88, no. 9, pp. 3149-3154. [https://doi.org/10.3168/jds.S0022-0322\(05\)72998-2](https://doi.org/10.3168/jds.S0022-0322(05)72998-2)
- Hosseinzadeh, S, & Dastmalchi-Saei, H 2014, 'Staphylococcal species associated with bovine mastitis in the North West of Iran: emerging of coagulase-negative staphylococci' *International Journal of Veterinary Science and Medicine*, vol. 2, no. 1, pp. 27-34. <https://doi.org/10.1016/j.ijvsm.2014.02.001>
- International Organization for Standardization 2002, *ISO 6579: Microbiology of food and animal feeding stuffs – Horizontal method for the detection of Salmonella spp*, ISO, Geneva
- Jamali, H, Radmehr, B, & Ismail, S 2014, 'Prevalence and antibiotic resistance of Staphylococcus aureus isolated from bovine clinical mastitis' *Journal of dairy science*, vol. 97, no. 4, pp. 2226-2230. <https://doi.org/10.3168/jds.2013-7509>
- Lüthje, P, & Schwarz, S 2006, 'Antimicrobial resistance of coagulase-negative staphylococci from bovine subclinical mastitis with particular reference to macrolide-lincosamide resistance phenotypes and genotypes' *Journal of Antimicrobial Chemotherapy*, vol. 57, no. 5, pp. 966-969. <https://doi.org/10.1093/jac/dkl061>
- McDougall, S, Hussein, H, & Petrovski, K 2014, 'Antimicrobial resistance in Staphylococcus aureus, Streptococcus uberis and Streptococcus dysgalactiae from dairy cows with mastitis' *New Zealand veterinary journal*, vol. 62, no. 2, pp. 68-76. <https://doi.org/10.1080/00480169.2013.84135>
- Minst, K, Märtilbauer, E, Miller, T, & Meyer, C 2012, 'Streptococcus species isolated from mastitis milk samples in Germany and their resistance to antimicrobial agents' *Journal of dairy science*, vol. 95, no. 12, pp. 6957-6962. <https://doi.org/10.3168/jds.2012-5852>
- Molina, D O 2010, *Análisis de la cadena de valor láctea de Honduras*, Pymerural/Pronagro, Tegucigalpa, Honduras
- Moon, J, S, Lee, A R, Kang, H M, Lee, E S, Joo, Y S, Park, Y H, ... & Koo, H C 2007, 'Antibiogram and coagulase diversity in staphylococcal enterotoxin-producing Staphylococcus aureus from bovine mastitis' *Journal of dairy science*, vol. 90, no. 4, pp. 1716-1724. <https://doi.org/10.3168/jds.2006-512>
- Oikonomou, G, Machado, V S, Santisteban, C, Schukken, Y H, & Bicalho, R C 2012, 'Microbial diversity of bovine mastitic milk as described by pyrosequencing of metagenomic 16S rDNA' *PloS one*, vol. 7, no. 10, pp. 1-14. <https://doi.org/10.1371/journal.pone.0047671>
- Patel, J B, Cockerill III, F, Eliopoulos, G, Jenkins, S, Lewis, J, Lim-bago, B, & Nicolau, D 2017, *M100S Performance standards for antimicrobial susceptibility testing*, Clinical and Laboratory Standards Institute. United States
- Pitkälä, A, Haveri, M, Pyörälä, S, Myllys, V, & Honkanen-Buzalski, T 2004, 'Bovine mastitis in Finland 2001—prevalence, distribution of bacteria, and antimicrobial resistance' *Journal of dairy science*, vol. 87, no. 8, pp. 2433-2441. [https://doi.org/10.3168/jds.S0022-0322\(04\)73366-4](https://doi.org/10.3168/jds.S0022-0322(04)73366-4)
- R. Core Team 2019, 'R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing' Vienna, Austria. <https://www.R-project.org/>
- Rall, V L M, Miranda, E S, Castilho, I G, Camargo, C H, Langoni, H, Guimarães, F F, ... & Júnior, A F 2014, 'Diversity of Staphylococcus species and prevalence of enterotoxin genes isolated from milk of healthy cows and cows with subclinical mastitis' *Journal of dairy science*, vol. 9, no. 2, pp. 829-837. <https://doi.org/10.3168/jds.2013-7226>
- Rato, M G, Bexiga, R, Florindo, C, Cavaco, L M, Vilela, C L, & Santos-Sanches, I 2013, 'Antimicrobial resistance and molecular epidemiology of streptococci from bovine mastitis' *Veterinary microbiology*, vol. 161, no. 3-4, pp. 286-294. <https://doi.org/10.1016/j.vetmic.2012.07.043>
- Roca, I, Akova, M, Baquero, F, Carlet, J, Cavalieri, M, Coenen, S, ... & Kahlmeter, G 2015, 'The global threat of antimicrobial resistance: science for intervention' *New microbes and new infections*, vol. 6, pp. 22-29. <https://doi.org/10.1016/j.nmni.2015.02.007>
- Saidi, R, Mimoune, N, Baazizi, R, Benaissa, M. H, Khelef, D, & Kaidi, R 2019, 'Antibiotic susceptibility of Staphylococci isolated from bovine mastitis in Algeria' *Journal of Advanced Veterinary and Animal Research*, vol. 6, no. 2, pp. 231. <https://doi.org/10.5455/javar.2019.f337>
- Sokal, R R, & Rohlf, F J 1995, *Biometry: the principles and practice of statistics in biological research*, Freeman and Company, New York: W. H.
- Sutra, L, & Poutrel, B 1994, 'Virulence factors involved in the pathogenesis of bovine intramammary infections due to Staphylococcus aureus' *Journal of Medical Microbiology*, vol. 40, no. 2, pp. 79-89. <https://doi.org/10.1099/00222615-40-2-79>
- Van Boeckel, T P, Brower, C, Gilbert, M, Grenfell, B T, Levin, S A, Robinson, T P, ... & Laxminarayan, R 2015, 'Global trends in antimicrobial use in food animals' *Proceedings of the National Academy of Sciences*, vol. 112, no. 18, pp. 5649-5654. <https://doi.org/10.1073/pnas.1503141112>
- Ventola, C L 2015, 'The antibiotic resistance crisis: part 1: causes and threats' *Pharmacy and Therapeutics*, vol. 40, no. 4, pp. 277.



Viridis, S, Scarano, C, Cossu, F, Spanu, V, Spanu, C, & De Santis, E P L 2010, 'Antibiotic resistance in Staphylococcus aureus and coagulase negative staphylococci isolated from goats with sub-


clinical mastitis' *Veterinary medicine international*, vol. 2010, id. 577060. <https://doi.org/10.4061/2010/517060>  
World Health Organization 2014, *Antimicrobial resistance: global report on surveillance*. World Health Organization, Geneva.



UNIVERSIDAD DE SAN CARLOS DE GUATEMALA  
FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA  
ESCUELA DE MEDICINA VETERINARIA


ANTIBIOTIC RESISTANCE AND SENSITIVITY PATTERNS OF  
BACTERIA ISOLATED FROM MASTITIS CASES IN DAIRY COWS  
IN HONDURAS


RESISTENCIA ANTIBIÓTICA Y SENSIBILIDAD EN AISLAMIENTO  
EN BACTERIAS EN MASTITIS EN VACAS LECHERAS


f.   
Br. Daniela Geovanna Martínez Matamoros



f. \_\_\_\_\_  
PhD. Dennis Guerra Centeno  
ASESOR PRINCIPAL

f.   
MSC. Manuel Lepe López  
ASESOR

  
f. \_\_\_\_\_  
Dr. Sc. Carlos Valdez Sandoval  
ASESOR

IMPRIMASE  
f.   
M.A. Rodolfo Chang Shum  
DECANO

